



# The Rajasthan Medical Journal

Indexed in Index Copernicus -( ICV- 2020- 50.77)  
Official Publication of the Medical and Health Department, Government of Rajasthan  
Published by S.M.S. Medical College, Jaipur

---

Volume : 16

ISSN : 0485-9561

Issue 4 : Oct 2022

---

## ORIGINALARTICLE

- Incidence and outcome of critically ill patients with COVID-19 and Acute kidney Injury in tertiary care centre of north-west India

## REVIEWARTICLE

- Overview of Materiovigilance Programme of India

## DRUG UPDATE

- Linzagolix

## CASE REPORT

- A Rare Case of Pulmonary Nocardiosis in Late Post Renal Transplant Period

## KNOWLEDGE CORNER

- Vaccination for Elderly: Easiest Way for Disease Prevention



# The Rajasthan Medical Journal

## Editor in-chief

Dr. Rajeev Bagarhatta, Principal and Controller

## Editors

Dr. Sudhir Mehta, Senior Professor (Medicine)

Dr. Amitabh Dube, Additional Principal

Dr. Monica Jain, Additional Principal

Dr. Rashim Kataria, Senior Professor (Neurosurgery)

## Associate Editors

Dr. Anita Singhal (Microbiology)

Dr. Kapil Gupta (Physiology)

Dr. Sunil Kumar Mahavar (Medicine)

Dr. Shruti Bhargava (Pathology)

Dr. Deepali Pathak (Forensic Medicine)

Dr. Laxmi Kant Goyal (Geriatric Medicine)

Dr. Rinki Hada (Physiology)

Dr. Shivankan Kakkar (Pharmacology)

Dr. Puneet Agarwal (Skin & V.D.)

Dr. Kopal Sharma (Pharmacology)

Dr. Vipin Garg (Biochemistry)

Dr. Nidhi Sharma (Pathology)

## Editorial Office

Medical Education Unit, S.M.S. Medical College, J.L.N. Marg, Jaipur – 302004

91-141-2518535. e-mail : medicaljournal.rajasthan@gmail.com

## Editorial Assistants

Manish Jain, Vinod Kumar Sharma

Published by S.M.S. Medical College, Jaipur, Rajasthan, printed at Government Press, Jaipur Rajasthan

While every effort is made by the editorial team, members of boards and the publishers to avoid inaccurate or misleading information appearing in the individual articles and additional, the responsibility for the same solely rests with the authors concerned. The RMJ, the publishers, and members of the editorial team and board accept no liability whatsoever, for the consequences of any such inaccurate or misleading data, opinion or statement.



# The Rajasthan Medical Journal

---

Volume : 16

ISSN : 0485-9561

Issue 4 : Oct 2022

---

	<i>Contents</i>	<i>Page No.</i>
ORIGINAL ARTICLE	Incidence and outcome of critically ill patients with COVID-19 and Acute kidney Injury in tertiary care centre of north-west India.....	01
REVIEW ARTICLE	Overview of Materiovigilance Programme of India.....	08
DRUG UPDATE	Linzagolix.....	16
CASE REPORT	A Rare Case of Pulmonary Nocardiosis in Late Post Renal Transplant Period.....	18
KNOWLEDGE CORNER	Vaccination for Elderly: Easiest Way for Disease Prevention.....	21
The RMJ Policy & Guidelines Instructions for Authors Renewal.....		23

## ORIGINAL ARTICLE

# Incidence and outcome of critically ill patients with COVID-19 and Acute kidney Injury in tertiary care centre of north-west India

Shashank Bhardwaj\*, Seetaram Singh\*, Tushar Gupta\*, Dhananjai Agrawal\*\*, Vinay Malhotra\*\*, Rakesh Gupta\*\*\*, Niranjana Gogoi\*, Gigin SV\*, Kavish Sharma\*, Shivi\*\*\*\*

### ABSTRACT

**Background:** The clinical manifestations of the COVID-19 range from asymptomatic to a fulminant and rapidly fatal infection. Studies show that the incidence of acute kidney injury (AKI) in COVID-19 patients vary widely from 0.5 to 80%.

**Objective:** This study was conducted to study the incidence and outcome of critically ill patients having COVID-19 infection along with AKI at a tertiary care center.

**Design:** A prospective cohort study done over a period of one month.

**Setting:** In this study, all adult ( $\geq 18$  years) patients admitted in Intensive Care Unit (ICU) and have tested positive for COVID-19 via RT-PCR test were included. All patients found to have AKI were assessed for comorbidities like diabetes, hypertension, Coronary artery disease, Stroke etc. Patients were followed up during the hospital stay.

**Results:** Incidence of AKI among COVID-19 patients was found to be 32.3%. There was significant ( $p < 0.05$ ) association of incidence of AKI with cough, fever, high total leucocyte count, use of face mask, use of high flow mask, diabetes and coronary artery disease (CAD). Cough ( $p = 0.01$ ), use of face mask ( $p = 0.006$ ), invasive mechanical ventilation ( $p = 0.0001$ ) and hypertension ( $p = 0.03$ ) were significantly associated with mortality among these patients. Mortality was highest among patients of AKI stage III (95.7%).

**Conclusion:** Incidence of acute kidney injury in critically ill patients with COVID-19 hospitalized in a tertiary care centre in western part of India was high. AKI

during hospitalization was associated with an increased risk of in-hospital death.

**Keywords:** COVID-19, Acute Kidney Injury, Critically ill

### INTRODUCTION

In December 2019, a series of cases of acute respiratory illness occurred in Wuhan, Hubei Province, China. This disease was identified as COVID-19, which is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. On March 11, 2020, the World Health Organization announced that the COVID-19 virus was officially a pandemic after it spread to more than 114 countries in just 3 months. More than 16 months after the first reported case, the virus is still spreading at a rapid rate with more than 100 million affected and more than 3 million dead. As India is struggling with a devastating second wave, the health care resources have been stretched to the limit. Physician, including nephrologist have had to rapidly adapt their practice to handle the large number of cases presenting to them. The clinical manifestations of the Covid-19 range from asymptomatic to a fulminant and rapidly fatal infection. The primary target of coronavirus disease is, the lungs but high incidences of acute kidney injury (AKI) have been reported. Initially, an incidence of AKI in hospitalized COVID-19 patients has been reported to be 34%<sup>1</sup>. In critically ill patients the incidence increased to 78%<sup>1</sup>. But, further studies found that the incidence of AKI in COVID-19 patients to vary widely from 0.5% to 80%<sup>2</sup>. This study was conducted to describe the clinical characteristics and risk factors for AKI and death among critically ill patients of COVID-19 who were treated in the ICU of our hospital.

\* Resident, \*\*Senior Professor, \*\*\*Assistant Professor, \*\*\*\*Senior Resident

Department of Nephrology, Sawai Man Singh Medical College and Hospital, Jaipur, Rajasthan

#### Corresponding Author:

Dr Shivi, Senior Resident, Department of Anaesthesia, Sawai Man Singh Medical College and Hospital, Jaipur, Rajasthan.

Contact no. 8860144064

Email id: Shashank.bhrdwaj@gmail.com



## MATERIALS AND METHODS

All patients presenting to the Intensive Care Unit (ICU) who tested positive for COVID-19 via RT-PCR were approached for inclusion into the study. Patient on immunosuppressive agents, with pre-existing chronic kidney disease and renal transplant recipients were excluded. After obtaining informed consent, a clinical profile was collected in a standard proforma. All patients found to have acute kidney injury were assessed for comorbidities like diabetes, hypertension, Coronary artery disease (CAD), stroke etc. Patients were followed up during the hospital stay and were assessed for need of oxygen (via face mask, high flow nasal cannula or high flow face mask), non-invasive or invasive mechanical ventilation. Proteinuria and hematuria were assessed

using a dipstick. Need for vasopressors/ inotropes was assessed. Stage of AKI was calculated using KDIGO guidelines. Renal replacement therapy in the form of hemodialysis or peritoneal dialysis was used according to standard practice guidelines.

## Definitions

AKI was identified according to the Guidelines of Kidney Disease: Improving Global Outcomes (KDIGO)<sup>3</sup>. It is defined as any of the following: an increase in Serum Creatinine of  $\geq 0.3$  mg/dL within 48 hours; an increase in Serum Creatinine of  $\geq 1.5$  times the baseline, which is known or presumed to have occurred within the past 7 days; or urine volume  $< 0.5$  mL/kg/hr for 6 hours.

Stage	Serum creatinine	Urine output
1	1.5–1.9 times baseline OR $\geq 0.3$ mg/dl ( $\geq 26.5$ $\mu$ mol/l) increase	$< 0.5$ ml/kg/h for 6–12 hours
2	2.0–2.9 times baseline	$< 0.5$ ml/kg/h for $\geq 12$ hours
3	3.0 times baseline OR Increase in serum creatinine to $\geq 4.0$ mg/dl ( $\geq 353.6$ $\mu$ mol/l) OR Initiation of renal replacement therapy OR, In patients $< 18$ years, decrease in eGFR to $< 35$ ml/min per $1.73$ m <sup>2</sup>	$< 0.3$ ml/kg/h for $\geq 24$ hours OR Anuria for $\geq 12$ hours

## RESULTS

The incidence of AKI was found to be 32.3%. Stage III AKI was among more than one third of patients (44.2%) followed by Stage I (28.8%) and Stage II (26.9%) Figure 1. About one third of patients were  $> 60$  years (34.2%). More than half of the patients were males (61.5%). The incidence of AKI was highest among patients of age  $> 60$  years (56.4%) and was lowest in  $< 30$  years (5%). The incidence of AKI was 24.54 times significantly higher among patients of age  $> 60$  years than  $< 30$  years (OR=24.54, 95%CI=3.06-196.50,  $p=0.003$ ). The incidence of AKI was higher among males (36.4%) than females (25.8%), however, the association was statistically insignificant ( $p>0.05$ ). Oxygen requirement

was the most common (92.5%) and dyspnoea was the second most common factor (73.3%). CVA was the least common factor (5.6%). The incidence of AKI was 6.42 times significantly higher among whom dyspnoea was present than absent (OR=6.42, 95%CI=2.15-19.16,  $p=0.0001$ ). There was significant ( $p<0.05$ ) association of incidence of AKI with cough, fever, face mask, high flow mask, diabetes and CAD (Table 1). The incidence of AKI was 5.44 times significantly higher among whom TLC (Total Leukocyte Count) was abnormal than normal (OR=5.44, 95%CI= 2.00-14.82,  $p=0.0001$ ). The factors found to be significant in univariate analysis were entered in the multivariate analysis. The multivariate analysis showed that the incidence of AKI was 12.44 times

significantly higher among patients of age >60 years than <30 years (Adjusted OR=12.44, 95%CI=1.42-127.76,  $p=0.03$ ) when adjusted for dyspnoea and high flow mask. Dyspnoea ( $p=0.01$ ) and high flow mask ( $p=0.0001$ ) was also significantly associated with the incidence of AKI in multivariate model. Cough ( $p=0.01$ ), face mask ( $p=0.006$ ), invasive mechanical ventilation ( $p=0.0001$ ) and hypertension ( $p=0.03$ ) were significantly associated with mortality (Table 2). Mortality was higher among patients of AKI stage III (95.7%) than stage I (80%) and stage II (42.9%) (Figure 2). The mortality was 82% significantly lower among patients of stage II than stage I (OR=0.18, 95%CI=0.03-0.97,  $p=0.04$ ). Among the patients who developed AKI, we found the incidence of 1+, 2+, 3+ proteinuria to be 40%, 38% and 5.8% respectively. 15.4% of the patients had no proteinuria. Interestingly no patient in our study had 4+ proteinuria on dipstick examination. Concurrent haematuria of 1+ and 2+ was seen in 46.2% and 1.9% of the patients, respectively.

## DISCUSSION

In this prospective cohort study conducted in Jaipur, Rajasthan, we observed an incidence of 32.3% of acute kidney injury in critically ill hospitalized patients with COVID-19. More than one third of these patients developed severe AKI, Stage III AKI. In our study, 14.28% of patients required renal replacement therapy.

The incidence of AKI reported in various studies varies from 0.5 to 80 percent<sup>2</sup>. In a meta-analysis<sup>4</sup> of approximately 13,000 mostly hospitalized patients, the incidence of AKI was 17 percent. Approximately 5 percent of patients required kidney replacement therapy (KRT) in this meta-analysis. The incidence of AKI seems to vary according to the number of critically ill patients included in the study. The high incidence of AKI in our study may be explained by the fact that we only included critically ill patients. The same may be the reason for higher number of patients requiring renal replacement therapy.

The incidence of AKI in critically ill patients, in pre COVID era has been found to be 39.3% in developed countries and 35.1% in the developing countries<sup>5</sup>. This is strikingly close to the incidence of AKI found in our study.

The incidence of AKI was higher among the 2600 patients who had COVID-19 compared with over 19,500 patients who were hospitalized for other reasons (31 versus 18 percent)<sup>6</sup>. This higher incidence of AKI could not be explained by differences in the traditional risk factors for AKI between the groups. COVID-19 remained associated with a higher rate of AKI despite controlling for demographic variables, comorbidities, frequency of hypotension, selected laboratory results (eg, complete blood count, baseline eGFR), and use of nephrotoxic medications, vasopressors, or mechanical ventilation. Whether AKI in COVID-19 patients portends worse mortality outcomes than non-covid patients requires further study.

The etiology of AKI in critically ill patients with COVID-19 appears to be multifactorial. In AKI-EPI study done in pre-covid era, etiology of AKI in descending order was as follows: sepsis, hypovolemia, drug related, cardiogenic shock, hepatorenal syndrome, and obstructive uropathy<sup>7</sup>. These are potential etiologies, even in critically ill patients of COVID-19 but whether SARS-CoV-2 causes a direct kidney infection remains controversial<sup>8</sup>. The role of “cytokine storm” and the high incidence of thrombosis in patients of COVID-19 remain understudied causes of AKI in this Cohort<sup>9,10</sup>.

A high incidence of AKI was seen in patients with respiratory failure. There was significant ( $p<0.05$ ) association of incidence of AKI with oxygen requirement with a face mask or a high flow mask. Incidence of AKI was also higher among patients who required Non Invasive and Invasive mechanical ventilation, but the incidence was not found to be statistically significant.

In other studies, the incidence of AKI was higher in patients on high oxygen support<sup>11</sup>.

Monitoring kidney function must therefore be emphasized even in patients with mild respiratory symptoms, and altered kidney function should be given particular attention after admission in clinical practice. Early detection and treatment of renal abnormalities, including adequate hemodynamic support and avoidance of nephrotoxic drugs, may help to improve the vital prognosis of COVID-19.

CT severity scores and markers of inflammation, such as C-reactive protein and ferritin, appeared to be

**Table 1: Distribution of various factors and its association with incidence of AKI**

Factors	No. (n=161)		With AKI		Without AKI		OR (95% CI)	P-value
	No.	%	No.	%	No.	%		
Dyspnea	118	73.3	48	40.3	70	59.7	6.42 (2.15-19.16)	0.0001*
Cough	72	44.7	15	20.8	57	79.2	0.37 (0.18-0.75)	0.005*
Fever	92	57.1	23	25	69	75	0.46 (0.23-0.90)	0.02*
Oxygen requirement	149	92.5	50	33.6	99	64.4	2.52 (0.53-11.96)	0.22
Nasal prongs	20	12.4	7	35.0	13	65.0	1.14 (0.42-3.07)	0.78
Face mask	28	17.4	1	3.6	27	96.4	0.06 (0.01-0.45)	0.0001*
High flow mask	22	13.7	14	63.6	8	36.4	4.65 (1.80-11.97)	0.001*
Non-invasive mechanical ventilation	52	32.3	18	34.6	34	65.4	1.16 (0.58-2.35)	0.66
Invasive mechanical ventilation	46	28.6	19	41.3	27	58.7	1.74 (0.85-3.56)	0.12
Diabetes	62	38.5	30	48.4	32	51.6	3.28 (1.65-6.52)	0.001*
Hypertension	63	39.1	26	41.3	37	58.7	1.94 (0.99-3.81)	0.06
CAD	17	10.6	10	58.8	7	41.2	3.46 (1.23-9.72)	0.01*
CVA	9	5.6	3	33.3	6	66.7	1.05 (0.25-4.37)	0.94
Need of vasopressors	48	29.8	19	39.6	29	60.4	1.58 (0.78-3.21)	0.19

**OR-Odds ratio, CI-Confidence interval, \*Significant****Table-2: Distribution of various factors and its association with mortality among critically ill patients.**

Factors	No. of patients	Expired		Alive		OR (95%CI)	p-value
		No.	%	No.	%		
Dyspnea	118	61	51.7	57	48.3	1.63 (0.80-3.32)	0.17
Cough	72	27	37.5	45	62.5	0.44 (0.23-0.84)	0.01*
Fever	92	42	45.7	50	54.3	0.77 (0.41-1.43)	0.41
Oxygen requirement	149	73	49.0	76	51.0	1.34 (0.40-4.42)	0.62
Nasal Prongs	20	7	35.0	13	65.0	0.53 (0.20-1.40)	0.19
Face Mask	28	7	25.0	21	75.0	0.29 (0.11-0.73)	0.006*
High flow mask	22	10	45.5	12	54.5	0.87 (0.35-2.14)	0.76
Non-invasive Mechanical	52	21	40.4	31	59.6	0.61 (0.31-1.20)	0.15

Ventilation							
Invasive Mechanical Ventilation	46	38	82.6	8	17.4	8.90 (3.79-20.91)	0.0001*
Diabetes	62	36	58.1	26	41.9	1.87 (0.98-3.57)	0.06
Hypertension	63	37	58.7	26	41.3	1.97 (1.04-3.76)	0.03*
CAD (Coronary Artery Disease)	17	10	58.8	7	41.2	1.59 (0.57-4.42)	0.36
CVA (Cerebrovascular Accident)	9	5	55.6	4	44.4	1.35 (0.35-5.23)	0.66
Need for Vasopressors	48	25	52.1	23	47.9	1.23 (0.62-2.42)	0.54
Renal Replacement Therapy	23	21	91.3	2	8.7	14.92 (3.36-66.17)	0.0001*
Haemodialysis	11	9	81.8	2	18.2	5.28 (1.10-25.27)	0.02*
Peritoneal dialysis	14	13	92.9	1	7.1	16.40 (2.09-128.65)	0.001*
AKI	52	40	76.9	12	23.1	6.22 (2.92-13.26)	0.0001*

OR-Odds ratio, CI-Confidence interval, \*Significant

higher among patients who had AKI compared with those who did not have AKI. However, the comparison between groups controlling for these inflammatory markers was not possible because the tests were not available for all the patients. A higher CT severity and marker of inflammation in those with AKI may be a pointer to a more severe disease, which may be the cause or effect or both in patients of COVID-19.

In our study, there was significant ( $p < 0.05$ ) association of incidence of AKI with older age, history of diabetes and CAD. There was a higher incidence of AKI among those with hypertension and history of CVA, but the incidence were not found to be statistically significant. Similarly, incidence of AKI was higher in those patients who required vasopressor support, but the incidence did not reach statistical significance. These are considered to be traditional risk factors for AKI. Patients with these comorbidities continue to be at risk for AKI, even among the patients of COVID-19 and frequent monitoring of renal functions of these patients may be warranted.

In our study, AKI was found to be significantly associated with mortality. Mortality was higher among patients of AKI stage III (95.7%) than stage I (80%) and stage II (42.9%). There was significant ( $p < 0.05$ ) association of mortality with requirement of renal replacement therapy. AKI has been found to be a risk factor for mortality in various studies of both COVID and non-COVID patients<sup>11,12</sup>. Whether AKI in COVID-19 patients portends worse mortality outcomes than non-COVID patients requires further study.

Among the patients who developed AKI, we found the incidence of 1+, 2+, 3+ proteinuria to be 40%, 38% and 5.8% respectively. 15.4 % of the patients had no proteinuria. Interestingly, no patient in our study had 4+ proteinuria on dipstick examination. Concurrent hematuria of 1+ and 2+ was seen in 46.2% and 1.9% respectively.

In most studies, where kidney biopsy was done in patients of COVID-19, the predominant kidney pathological finding was Acute Tubular Necrosis (ATN), but glomerular lesions have been reported in a minority of patients<sup>13-17</sup>. These findings explain the urinalysis findings in our patients. A renal biopsy could not be carried out in patients in our centre due to resource limitations, and the fact that all patients enrolled in the study were critically ill and many were haemodynamically unstable.

Even though this study included a substantial number of patients from a tertiary care hospital, it has several limitations. First, a baseline serum creatinine was not available for all patients, which may have led to an underestimation/overestimation of AKI or erroneous associations. Second, clinical data of patients after discharge were lacking, so we could not assess COVID-19 effects on long-term outcomes. The precise impact of COVID-19 on kidney structure and function and the incidence of chronic kidney disease in these patients require further investigation.

Our study represents the reality of public hospitals in our country. The findings described here, can help, not only in planning of future studies but also in




Figure 1: Incidence of AKI in COVID-19 patients according to AKI stages.

future planning of ICU care for critically ill patients of COVID-19. A high number of critically ill patients required renal replacement therapy and provisions for the same are an essential part of care for critically ill patients. The facility for the same is lacking in large parts of the country, and requires urgent attention.

In conclusion, the incidence of acute kidney injury in critically ill patients with COVID-19 hospitalized in Jaipur, was high. AKI during hospitalization was associated with an increased risk of in-hospital death. Clinicians should increase their awareness of kidney disease in hospitalized patients with COVID-19. Early detection and effective intervention of kidney involvement may help to reduce deaths of patients with COVID-19.

REFERENCES

1. Argenziano MG, Bruce SL, Slater CL. et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020; 369:m1996.

2. Jia H. Ng, Vanesa Bijol, Matthew A. Sparks et al. Pathophysiology and pathology of acute kidney injury in patients with COVID-19. *Advances in Chronic Kidney Disease* 2020; 27(5):365-376.

3. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl.* 2012;2(1):1–138.

4. Juarez SY, Qian L, King KL et al. Outcomes for Patients With COVID-19 and Acute Kidney Injury: A Systematic Review and Meta-Analysis. *Kidney Int Rep.* 2020;5(8):1149. Epub 2020 Jun 25

5. Melo FAF, Macedo E, Fonseca Bezerra AC et al. A systematic review and meta-analysis of acute kidney




Figure 2: Mortality according to stages of AKI in COVID-19 patients

injury in the intensive care units of developed and developing countries. *PLoS One.* 2020 Jan 17;15(1):e0226325.

6. Moledina DG, Simonov M, Yamamoto Y et al. The Association of COVID-19 With Acute Kidney Injury Independent of Severity of Illness: A Multicenter Cohort Study. *Am J Kidney Dis.* 2021;77(4):490. Epub 2021 Jan 8.

7. Hoste EA, Bagshaw SM, Bellomo R et al. Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med.* 2015 Aug;41(8):1411-23.

8. Lite C, Ahmed SSSJ, Juliet M et al. SARS-CoV-2/human interactome reveals ACE2 locus crosstalk with the immune regulatory network in the host. *Pathog Dis.* 2021;79(2)

9. Minami, T., Iwata, Y. & Wada, T. Renal complications in coronavirus disease 2019: a systematic review. *InflammRegener* 40,31 (2020).

10. Ahmadian E, HosseiniyanKhatibi SM, RaziSoofiyan S et al. Covid-19 and kidney injury: Pathophysiology and molecular mechanisms. *Rev Med Virol.* 2021 May;31(3):e2176.

11. Bowe B, Cai M, Xie Y et al. Acute Kidney Injury in a National Cohort of Hospitalized US Veterans with COVID-19. *Clin J Am SocNephrol.* 2020;16(1):14. Epub 2020 Nov 16.

12. Hirsch JS, Ng JH, Ross DW et al. Acute kidney injury in patients hospitalized with COVID-19. *Kidney Int.* 2020;98(1):209-218. Epub 2020 May 16.

13. Kudose S, Batal I, Santoriello D et al. Kidney Biopsy Findings in Patients with COVID-19. *J Am SocNephrol.* 2020;31(9):1959. Epub 2020 Jul 17.



14. Golmai P, Larsen CP, DeVita MV et al. Histopathologic and Ultrastructural Findings in Postmortem Kidney Biopsy Material in 12 Patients with AKI and COVID-19. *J Am Soc Nephrol.* 2020;31(9):1944. Epub 2020 Jul 16.
15. Werion A, Belkhir L, Perrot M et al. SARS-CoV-2 causes a specific dysfunction of the kidney proximal tubule. *Kidney Int.* 2020;98(5):1296. Epub 2020 Aug 10.
16. Akilesh S, Nast CC, Yamashita M et al. MulticenterClinicopathologic Correlation of Kidney Biopsies Performed in COVID-19 Patients Presenting With Acute Kidney Injury or Proteinuria. *Am J Kidney Dis.* 2021;77(1):82. Epub 2020 Oct 10.
17. Farkash EA, Wilson AM, Jentzen JM. Ultrastructural Evidence for Direct Renal Infection with SARS-CoV-2. *Am Soc Nephrol.* 2020;31(8):1683. Epub 2020 May 5.

## REVIEW ARTICLE

### Overview of Materiovigilance Programme of India

Arun Singh\*, Monica Jain\*\*, Rupa Kapadia\*\*, Uma Advani\*\*\*, Dharendra Kumar Mahawar\*\*\*\*, Shivankan Kakkar\*\*\*\*, Jaya Dadhich \*\*\*\*\*

#### ABSTRACT

The Medical devices vigilance also known as Materiovigilance, is the collection, assessment, reporting and identification of trends in incidents resulting from the uses of medical devices (ex. instrument, apparatus, implement, machine, appliance, implant, reagent for in vitro use, software, material or other similar or related article etc.). To ensure the safety of medical devices in India, a Materiovigilance Programme of India (MvPI) has been implemented from government of India in 2013 and it was formally launched on July 6, 2015, with collaboration of Sree Chitra Tirunal Institute for Medical Sciences and Technology (SCTIMST) as the National Coordinating Center (NCC). The collected and generated safety data will support to Central Drug Standard Control Organisation (CDSCO) for giving recommendation for safe use of medical devices in Indian population.

**Keywords:** Central Drug Standard Control Organisation (CDSCO), Materiovigilance, Medical Device, Safety.

#### INTRODUCTION

Materiovigilance is the supervision of adverse events occurring due to use of medical devices in patients. It is helping to develop the healthcare system of India. The word materiovigilance is derived from two Greek words "materio"- which means materials and "vigilare" meaning to keep awake or alert, to keep watch. Materiovigilance, as defined by the World Health Organization (WHO), is the science and activities relating to the detection, assessment, understanding and prevention of adverse events caused by medical devices. The purpose of medical devices for the identification, treatment,

diagnosis, and cure of illness that may sometimes initiate adverse reactions which needs to be monitored and prevented. So, thus a vigilance system is required to supervise the adverse events<sup>1</sup>. In the beginning, the marketed medical devices were regulated according to the Drugs and Cosmetics Act, 1940 and Rules 1945. After that in year 2017, an absolute rule was outlined to regulate the medical devices that existing in the Indian market called as Medical Device Rules, 2017<sup>2</sup>.

Medical device includes any instrument, implant, machine, appliances, equipment, grafting material, reagent for in vitro use or other article used on itself or combinedly as well as software needed for it to function properly, which is proposed to be used on patients by manufactures for the following scopes<sup>3</sup>:

- (a) For diagnostic, prevention, control, treating or lessening the diseases.
- (b) For diagnostic, control, treating, for reducing or counteracting an injury, impairment, and disabilities.
- (c) For learning, substituting, or altering component of the anatomy or a physiological process.
- (d) For grand achievement in conception.

#### MATERIOVIGILANCE PROGRAMME OF INDIA

In current scenario, there are need to regulate the safety surveillance of medical devices in India for prevention of unexpected events due to uses of medical device. The Ministry of Health and Family Welfare (MOHFW) has authorized the MvPI, an organisation for monitoring the safety of medical equipment affiliated from the Government of India in year 2013. The MvPI was officially introduced on July 6, 2015 at Indian

---

\*Senior Resident, Department of Pharmacology, SMS Medical College, Jaipur

\*\*Senior Professor, Department of Pharmacology, SMS Medical College, Jaipur

\*\*\*Associate Professor, Department of Pharmacology, SMS Medical College, Jaipur

\*\*\*\*Assistant Professor, Department of Pharmacology, SMS Medical College, Jaipur

\*\*\*\*\*Senior Demonstrator, Department of Pharmacology, SMS Medical College, Jaipur

Corresponding Author

Dr Arun Singh

Pharmacopoeia Commission (IPC), Ghaziabad by the Drugs Controller general India (DCGI). All prescriber, nursing officers and patients/consumers be made able to report Medical Devices Associated Adverse Events (MDAEs) to SreeChitraTirunal Institute of medical Sciences & technology (SCTIMST), Thiruvananthapuram or National Coordination Centre (NCC) - IPC<sup>4</sup>. The reported, collected and generated safety data will support to make easier in giving recommendations to Central Drugs Standard Control Organisation (CDSCO) and on directing of safe use of medical devices in Indian Population. Due to increasing number of cases of malfunctioning in many medical equipments and numerous medical device related adverse events, developing to further complications in patients and in most severe cases life-threatening conditions, the significance and necessity of MvPI are rising from previously days. Soon after in year 2018 onward, the IPC functions as NCC for MvPI as well NCC for Pharmacovigilance Programme in India (PvPI). MvPI has Medical Device Adverse Events Monitoring Centres (MDMCs) all over the country. To make sure effective of Adverse Events reporting from all respective individuals centres, this programme had established several tools for Adverse Event reporting to create India specific data<sup>5</sup>. For implementation of MvPI effectiveness, it is recognized that the integration of biomedical engineering department (BMED) at the health facilities/institutions is fundamental for further harmonisation with other departments. Hence, for functions as MDMCs, the

preferences have been given to the institutes with BMED.

### Different Medical Device Adverse Events Monitoring Centres Across India:

In India, there are ten MDMCs which have been recognized under MvPI that work to report MDAEs whose counts has been expanded to 150. Since 2015, the adverse events reporting rate has also increased. Through MvPI, more than 7000 reports have been submitted to IPC, Ghaziabad. For appropriate identification, reporting and collection of any suspected or definite MDAEs, MDMCs are responsible. MDAEs are classified into five categories: not related, unlikely, possible, probable, and causal relationship, respectively. MDMCs will send the reported data to NCC-IPC every month for review and analysis. IPC, Ghaziabad is the individual custodian of MvPI database. NCC is responsible to coordinate with all MDMCs nationwide and connects all concerned issues to the CDSCO. Apart from this, they collaborate with global authorities. It also provides financial support to SCTIMST, the National Health Systems Resource Centre (NHSRC), and MDMCs. SCTIMST functions as the NCC and gives help in all technical matters. NHSRC works as a technical support partner in the MvPI. Ultimately, all concerns are transferred to the CDSCO, i.e., a national regulatory authority ensuring safety. CDSCO is responsible for taking decisions on recommendations by the NCC-MvPI. The organizational structure of MvPI is represented in Figure 1.




Figure 1: Diagrammatic Representation of MDEM Under MvPI



## OBJECTIVES OF MATERIOVIGILANCE PROGRAM OF INDIA

The MvPI was began with the objectives to make safe the healthcare system and make sure the safety of medical device users and others by decreasing the recurrences of adverse events and malfunctions of medical equipments<sup>6</sup>.

The objectives are:

- (a) To generate a countrywide system for monitoring of patient safety
- (b) To scrutinise the risk–benefit ratio of medical equipment uses
- (c) To create evidence-based data on the safety of medical equipment
- (d) To support CDSCO in making precise decision process on the use of medical equipment
- (e) To convey the safety information on the use of medical equipment among stakeholders to diminish the risk
- (f) To develop as a national coordination center of excellence for materiovigilance activities
- (g) To work together with other healthcare organizations and international body for the exchange of data information and its management

## Regulation of Medical Devices Under Drug and Cosmetic Act:

In our country, at present days only notified medical devices are regulated as Drugs under the Drugs and Cosmetics Act 1940 and Rules made there under in 1945 that are given as following<sup>7</sup>: (A) Substances used for in vitro diagnosis and surgical dressings, surgical bandages, surgical staples, surgical sutures, ligatures, blood and blood component collection bag with or without anticoagulant covered under sub-clause (i); (B) Substances including mechanical contraceptives (condoms, intrauterine devices, tubal rings), disinfectants and insecticides notified under subclause (ii); and (C) Devices notified from time to time under subclause (iv), of clause (b) of section 3 of the Drugs and Cosmetics Act, 1940

## POST MARKETING SURVEILLANCE APPROACHES IN WORLDWIDE

In 2011, a global organization with the designation of International Medical Device Regulators

Forum (IMDF) was established with the primary goal of accelerating the coordination of international medical devices regulation and to reinforce the foundation of Global Harmonization Task Forces (GHTF). Ten countries participated in IMDF management committee i.e., Australia, Brazil, Canada, China, European Union, Japan, Russia, Singapore, South Korea, and USA. All these countries have founded their individual surveillance systems for active and passive monitoring of MDAEs. The United State of America (USA) has founded a universal recognized system known as Food and Drug Administration (FDA) for the monitoring of food, medications, vaccines, and medical devices. In the same country, a mandatory and voluntary two schemes of medical device reporting have also been proposed<sup>8</sup>, whereas United Kingdom (UK) has "Vigilance Reporting Scheme" as well the "Adverse Event Scheme" for the post-marketing surveillance of medical devices<sup>9</sup>. The medical device manufacturers must take Vigilance Reporting Scheme and they should report adverse events within prescribed timeline. Although, it is voluntary for the healthcare professionals, hospital engineers and patients / consumers to take Adverse Event Scheme and report immediately. This scenario is different in Europe as all medical devices related issues are handed directly through National Competency Authority (NCA) from the manufacturer, while it is compulsory for general prescribers, doctors, and nurses to report the adverse event to manufacturer as well as NCA. The manufacturers are needed to submit a preliminary report of the serious medical device related adverse event within 2 calendar days. Later this, if the manufacturer realizes any relation between a medical device and life threatening or death or any health hazard, the appraisal is required to be reviewed within 10 days. Apart from this, very less serious or non-serious cases can be reported within 30 calendar days<sup>10</sup>. In Japan country, medical devices are being regulated by the Pharmaceutical and Medical Device Agency that approaches come under certain rules and regulations which are required to be fulfilled for certification, quality assurance, and licensing including Japanese medical products<sup>11</sup>. In France, the medical device related adverse events are regulated by the "National Surveillance Commission" which is also known as the "Commission of Materiovigilance". In Australia, the regulatory body, "Therapeutic Goods Administration" come under the Department of Health of the Australian Government that

regulates the medical device related adverse events<sup>12</sup>. Therapeutic Goods Administration keep data upto 5 years. In Canada, “Health Canada” Regulatory Authority regulates and ensure that all medical devices are safe and rationally used in both phases: pre- and post marketing surveillance and should be reported within 10 days after the manufacturer of a medical device becomes aware of

an incident. In India, as already discussed that “Materiovigilance Programme of India” has been regulated the medical device related adverse events and reported to IPC within 15 calendar days of becoming aware of an event and for non-serious events reporting to be done within 30 calendar days.

### Pharmacovigilance Vs Materiovigilance

Materiovigilance is differ from Pharmacovigilance as given following:

	Pharmacovigilance	Materiovigilance
1. Definition	Pharmacovigilance is a system of monitoring Safety and Effectiveness of Drugs and other Pharmaceuticals.	Materiovigilance is a system of Monitoring Safety and Effectiveness of a Medical Device.
2. Programme Name in India	Pharmacovigilance Program of India	Materiovigilance Program of India
3. Launched Programme	In 2010	In 2015
4. Location of National Coordination Centre	Indian Pharmacopoeia Commission, Ghaziabad, previously in AIIMS, Delhi.	Indian Pharmacopoeia Commission, Ghaziabad, SCTIMST and NHSRC.
5. Regulation	Finally, CDSCO will take necessary action against ADRs.	Finally, CDSCO will take necessary action against MDAEs.
6. Numbers of centres	At present, India have 395 adverse drug reaction monitoring centres in various medical colleges.	In India, currently medical device adverse events monitoring centres number has been increased to 150.

### Reporting of Medical Device associated Adverse Events (MDAEs)

#### What to Report

All types of suspected Medical Device related Adverse Events (MDAEs) can be reported whether they are serious or non-serious, frequent, or rare regardless of an established causal relationship. A Medical device can be any material, implant, incubator, instrument, apparatus, reagent, and any diagnostic tests, etc. Adverse Events description, Details of adverse event including description of device (deficiency or malfunction), clarification of threats associated with that device and the

associated risk of patients with previous use can be provided in the MDAEs reporting form.

#### Who can Report

Practitioners, Biomedical Engineers, Clinical Engineers, Hospital Technology Managers, Pharmacists, Nurses, Technicians can report medical device adverse events.

#### Why to Report?

All associated medical device related adverse events should be reported in order to protect public health. Any unreported adverse events may lead to serious or death of patients.

## Overview of Materiovigilance Programme of India

### Where to Report

The Healthcare workers (doctors, nurses, dentists, pharmacists), Manufactures and Patient/Consumers can report MDAEs to SCTIMST or NCC. Duly filled MDAEs Reporting Form can be send to SCTIMST or NCC- MvPI, Biomedical Technology Wing, Poojappura, Thiruvananthapuram, 695012, Kerala, India or Can directly email with scanned copy of the duly filled form to mvpi@sctimst.ac.in.

### How to Report

MDAEs Medical Device Adverse Event Reporting Form [Figure 2] can be downloaded from the website of IPC (www.ipc.gov.in) to report adverse event associated with medical devices. MDAEs can also be reported via PvPI helpline number (1800 180 3024) on weekdays from 9:00 am to 5:30 pm.



### MEDICAL DEVICE ADVERSE EVENT REPORTING FORM

Materiovigilance Programme of India

FOR MDMC/NCC USE ONLY												
Type of report: Initial <input type="checkbox"/> Follow-up <input type="checkbox"/>				Report No. _____								
<b>A. PATIENT DETAILS</b>												
1. Patient Hospital ID _____				3. Age at time of Event or Date of Birth _____								
2. Sex: M <input type="checkbox"/> F <input type="checkbox"/>				4. Weight (Kg) _____								
<b>B. EVENT DETAILS</b>												
1. Event description-  												
Reason for the Event(Tick) a) Electrical <input type="checkbox"/> b) Mechanical <input type="checkbox"/> c) Electronic <input type="checkbox"/> d) Biocompatibility <input type="checkbox"/> e) Clinical application error <input type="checkbox"/>												
2. Severity of the event (Yes <input type="checkbox"/> No <input type="checkbox"/> if yes please specify following <input type="checkbox"/> Death (_____) <input type="checkbox"/> cause congenital-anomaly <input type="checkbox"/> Life threatening <input type="checkbox"/> Required intervention to prevent death or impairment of body function <input type="checkbox"/> Hospitalization/Prolonged impairment/damage <input type="checkbox"/> Disability <input type="checkbox"/> Other (specify) _____												
3. Date of event - (dd/mm/yyyy)												
4. Location of the event- OPD <input type="checkbox"/> IPD <input type="checkbox"/> Others (Please specify) _____												
5. Device category: (A) Therapeutic <input type="checkbox"/> Diagnostics <input type="checkbox"/> Both <input type="checkbox"/> (B) Implantable device <input type="checkbox"/> Non Implantable device <input type="checkbox"/> (C) Single use device <input type="checkbox"/> Reusable device <input type="checkbox"/> Reuse of manufacture marked single use device <input type="checkbox"/>												
6. Date- <table border="1" style="width: 100%;"> <tr> <td>Last preventive maintenance</td> <td>Last calibration</td> </tr> <tr> <td> </td> <td> </td> </tr> </table>									Last preventive maintenance	Last calibration		
Last preventive maintenance	Last calibration											
7. Location of device after the incident: Place of use <input type="checkbox"/> Place of reporter <input type="checkbox"/> Place of Manufacture/vendor <input type="checkbox"/> With patient or end user <input type="checkbox"/>												
8. Is device in use after incident Yes <input type="checkbox"/> No <input type="checkbox"/>												
9. (A) Is same model device available in organisation? Yes <input type="checkbox"/> No <input type="checkbox"/> If yes, Quantity _____ (B) Organization - Healthcare facility <input type="checkbox"/> Manufacturer <input type="checkbox"/>												
<b>C. MEDICAL DEVICE(S) DETAIL</b>												
Name of Medical Device (1)	Manufacturer (2)	Brand Name (3)	Model No. (4)	Serial No. (5)	Batch No./ Lot No. (6)	Catalogue No. (for instruments only)	Date of installation/ implantation/ explanation (8)	List of Accessories (9)				
10. Actions taken immediately after incident				11. A Whether other medical devices were being used at same time with above device for therapeutic or diagnostic service? If yes, please specify _____								
				11. B. Any history of adverse event(s) from device with same serial/model/catalogue number. If yes please specify _____								

D. REGULATORY DETAILS			E. REPORTER DETAILS of MvPI CENTRE
Manufacturer name:	Entity legally representing the Manufacture:	Notified Body name in:	Name and Professional Address: _____
Regulator in Country of origin:	Country:	(i) Country of Manufacturing :	Pin: _____ E-mail _____
Regulatory status in origin country:		(ii) In India:	Tel. No. (with STD code) _____
			Designation: _____
			Signature: _____
			Date of this report ____/____/____
<b>F. Causality Assessment Details</b> Completed <input type="checkbox"/> In Progress <input type="checkbox"/> Awaited <input type="checkbox"/>			
Additional Information:			
<small>Confidentiality: The patient's identity is held in strict confidence and protected to the fullest extent. Programme staff is not expected to and will not disclose the reporter's identity in response to a request from the public. Submission of a report does not constitute an admission that medical personnel or manufacturer or the product caused or contributed to the adverse event.</small>			



**National Collaborating centre-Materiovigilance Programme of India.**

**Sree Chitra Tirunal Institute for Medical Sciences and Technology (SCTIMST)** under the Department of Science & Technology, Government of India. Biomedical Technology Wing, Poojappura, Thrivananthapuram 695012, Kerala. Phone: 91- 471 – 2340411, Fax: 91- 471 -2341814, Email: [head-bmtw@sctimst.ac.in](mailto:head-bmtw@sctimst.ac.in).



**National Coordination Centre-Materiovigilance Programme of India.**

**Indian Pharmacopoeia Commission (IPC)**, Ministry of Health and Family Welfare, Government of India, Sector-23, Rajnagar, Ghaziabad-20002, Tel.: 0120-2783400, 2783401, and 2783392, FAX: 0120-2783311, Email: [ipclab@vsnl.net](mailto:ipclab@vsnl.net), [pvpi.ipcindia@gmail.com](mailto:pvpi.ipcindia@gmail.com)



**Technical support and Resource Centre- Materiovigilance Programme of India.**

**National Health System Resource Centre (NHSRC)**, NIHF campus Baba Gangnath marg, Munirka, New Delhi-110067, Phones: 011 26108982 / 83 / 84 / 92 / 93, Fax: 011-26108994 Email: [nhsr.india@gmail.com](mailto:nhsr.india@gmail.com).

**Where to report**

- Duly filled Medical Device Adverse Event Reporting Form can be send to Sree Chitra Tirunal Institute for Medical Sciences and Technology (SCTIMST), National Collaboration Centre-Materiovigilance Programme of India, Biomedical Technology Wing, Poojappura, Thrivananthapuram 695012, Kerala, India.
- Or Can directly email the duly filled form to [mvpi@sctimst.ac.in](mailto:mvpi@sctimst.ac.in).
- Call on Helpline no. 1800 180 3024 to report Adverse event.

**Event description** Details of adverse event including description of device (deficiency or malfunction), clarification of hazards associated with device and the associated risk of patient, user or person any possible risk to patient associated with previous use.

**Additional Information** Other relevant information related to treatment should be provided.

Figure 2: Medical Device Adverse Reporting Form

### Example for Materiovigilance:

The manufacturer has issued a batch of blood glucose test strips. According to instructions the patients use the strips. Because of wrong readings, wrong insulin dosage was taken by the patient which leads to hypoglycaemic shock and hospitalization.

Another example is the patients facing problems related to malfunction of cardiac pacemaker that leads to output failure or failure to capture or pacemaker-medicated tachycardia or runaway pacemaker etc. Because of these problems patients can suffer syncope and cardiac arrest.

### Future Prospects of Materiovigilance Programme of India

If the MvPI has to be successful, the contribution at root level has to be accelerated. So, there is need to provide awareness among health-care workers and the general public about MvPI has to be strengthened. For increasing the awareness, we should know what exactly to report, that is potential barriers in reporting MDAEs. If they reported to regulatory authorities timely, it will help to those diseased patients that will be affected due to deficiency and malfunctions of using medical devices. Academic curricula MBBS, Postgraduate programs, nursing, dental and paramedical courses should be updated to include MvPI, and the knowledge of reporting MDAEs. The number of materiovigilance centres are increasing continuously for implementation of MvPI. MvPI has key role in prevention of medical device associated health hazard.

### CONCLUSION

In a few past years, the medical device's uses are found to be frequently by doctors across all over world including India. In spite of that, there are not satisfactory tool to safe guard the patients from the undesired events. Materiovigilance Programme of India is a good initiative to make sure the safety of medical device. This programme will be also significantly reducing the risk associated to the use of medical devices by avoiding the repetition of aversive effects. Therefore, it is the moral responsibility for all healthcare professionals to report MDMEs to IPC, Ghaziabad through submission of reporting form so that CDSCO can give recommendations to prevent medical device related health hazards.

### REFERENCES

1. Johnson JA. FDA Regulation of Medical Devices. CRS Report for Congress Prepared for Members and Committees of Congress; 25 June 2012. Available from: <http://www.fas.org/sgp/crs/misc/R42130.pdf>. [Cited on October 7,2022]
2. 2.Medical Device Rules 2017.pdf [Internet]. Available from: <https://www.dfda.goa.gov.in/attachments/article/419/MedicalDevice%20Rules%202017.pdf>. [Cited on October 7,2022].
3. Dr. John G. Webster, Professor of Biomedical Engineering at the University of Wisconsin: Encyclopaedia of Medical devices and Instrumentation, Second edition, John Wiley & Sons: 2010.
4. Available from: <http://www.Users/Aiims%2044/Desktop/materiovigilance/materiovigilance%20programme%20of%20India.pdf>. [Last accessed on October 8,2022].
5. Indian pharmacopoeia commission, available at <https://www.ipc.gov.in>. [Last accessed on October 8,2022].
6. Available from: <http://www.Users/Aiims%2044/Desktop/materiovigilance/materiovigilance%20programme%20of%20India.pdf>. [Last accessed on October 8,2022]
7. Central Drug Standard Control Organization. Available from: <https://cdsco.gov.in/opencms/opencms/en/Medical-Device-Diagnostics/Medical-Device-Diagnostics/>. [Last accessed on October 8,2022].
8. Volume : VII, Issue : VI, June – 2018; DOI: 10.36106/ijsr. 2. Materiovigilance in EU, available at [https://www.lambda\\_cro.com/materiovigilance-in-eu/](https://www.lambda_cro.com/materiovigilance-in-eu/).
9. Manu M, Anand G. A review of medical device regulations in India, comparison with European Union and way-ahead. *Perspect Clin Res* 2022;13:3-11.
10. EU MDR Vigilance Reporting Requirements and MEDDEV 2.12-1 Rev 8: What Has Changed?



11. Japan MHLW & PMDA Medical Device and Pharmaceutical Regulations. Available from: <https://www.pacificbridgemedical.com/regulation/Japan-medical-device-pharmaceuticalregulations/#:~:text=Current%20Japan%20PMDA%20regulations%20are,Gen%20Therapy%20Products%2C%20and%20Cosmetics> [Last accessed on 2022, Oct 10].
12. Gupta P, Janodia MD, Jagadish PC, Udupa N. Medical device vigilance systems: India, US, UK, and Australia. *Med Devices (Auckl)* 2010;3:67-79.

## DRUG UPDATE

### Linzagolix

Monica Jain\*, ShivankanKakkar\*\*, Yatendra Singh\*\*\*


Linzagolix has been recently approved for the management of moderate-to-severe symptoms associated with leiomyomas (in the reproductive age group females) in the European Union (EU), whereas it is under regulatory review status in the United States of America (USA). Linzagolix is awaiting its entry in the Indian markets.

In the reproductive age group females, there are Oestrogen-related disorders like fibroids (uterine leiomyomas), endometriosis (endometrium-like tissue outside the uterine cavity) and adenomyosis (pathologically endometrial glands and stroma in uterine wall muscle).

Leiomyomas symptomatically present as- heavy menstrual bleeding (can result in anaemia), dysmenorrhea, pelvic pain, frequent urination, dyspareunia and it also increases risk for pregnancy complications, which are mainly managed surgically (mainly hysterectomy), Pharmacological management options available are like Gonadotropin releasing hormone (GnRH) agonists, Progestin-releasing IUD and Tranexamic acid. In recent studies, a promising alternative has emerged that is Linzagolix. It has been found to effectively improve symptoms with significant reduction in uterine volume.<sup>1</sup>

Chemical Name

3 - { 5 - [ ( 2 , 3 - d i f l u o r o - 6 - methoxyphenyl)methoxyl]-2- fluoro-4- methoxyphenyl } -2,4- dioxo-1,2,3,4- tetrahydrothieno [3,4-d] pyrimidine-5- carboxylic acid



Linzagolix is a selective, non-peptide, orally active Gonadotropin releasing hormone (GnRH) antagonist, indicated for treatment of moderate-to-severe symptoms related with leiomyomas in reproductive age group females<sup>2</sup>.

#### Preclinical data

In animal studies on ovariectomized cynomolgus monkey models reduction in serum LH levels at 8 hours was found which sustained for over 24 hours. Whereas, in intact female cynomolgus monkeys, dose -dependent partial or complete blockage of GnRHsignalling was seen, with resumption of normal menstrual cycles and hormonal secretion within one cycle on withdrawal<sup>3</sup>.

#### Clinical trials data

In phase 1 trials, it was found that Linzagolix can be orally administered (taken with or without food) and gets rapidly absorbed. It is highly bound to plasma proteins, with a half-life of 15 hours. It is mainly excreted in urine, then in faeces, majorly unchanged<sup>3</sup>.

Linzagolix in phase 2 and phase 3 trials effectively reduced pain and other symptoms and uterine and fibroid size as a selective GnRH receptor antagonist, blocking the hypothalamic pituitary-gonadal axis, resulting in dose dependent reduction in serum levels of luteinising hormone (LH), follicle-stimulating hormone (FSH) and estradiol. The GnRH receptors are coupled with Gq/11 which on activation increases intracellular Ca<sup>2+</sup> flux, Linzagolix dose-dependently inhibits GnRH-stimulated Ca<sup>2+</sup> flux.

Linzagolix improved dysmenorrhea and non-menstrual pelvic pain in the randomised, double-blind, placebo-controlled phase 3 conducted in Europe and the USA in women with moderate-to-severe endometriosis-associated pain<sup>3</sup>.

\*Senior professor, Department of pharmacology, SMS medical college

\*\*Assistant professor, Department of pharmacology, SMS medical college

\*\*\*First year resident, Department of pharmacology, SMS medical college

#### Corresponding Author:

Dr. Yatendra Singh

First year resident, Department of Pharmacology, SMS medical college

E-mail : singh.yatendra1801@gmail.com

Treatment with high-dose linzagolix 200 mg once daily for 12 weeks followed by a 100 mg/day maintenance dosage for a further 12 weeks significantly reduced uterine volume and adenomyosis-related symptoms, including pelvic pain and dysmenorrhea, in an exploratory phase 2 trial in 8 premenopausal women with symptomatic uterine adenomyosis<sup>3</sup>.

In clinical studies, the most common adverse events associated with Linzagolix were hot flushes and headache, more commonly at higher doses. These events were less frequent with Linzagolix with add-back therapy.

In phase 3 trials, less than 6 month use of Linzagolix 200 mg daily, reduction in size of leiomyomas was found, which may increase in size on cessation of drug. It should not be given more than 6 months due to risk of reduction of Bone Mineral Density on long term use<sup>3</sup>.

In phase 3 trials, high lipid levels (LDL cholesterol, HDL cholesterol and triglycerides) were seen, which generally were higher in Linzagolix without add-back therapy (ABR). These levels reduced after the treatment but had not reached the baseline levels<sup>3</sup>.

It should be given cautiously to patient with a history of depression and/or suicidal ideation as mood disorders like emotional lability, alteration in mood and depression are seen with use of GnRH antagonists including Linzagolix<sup>3</sup>.

Relatively contraindicated for patients with moderate or severe kidney impairment or with end stage kidney disease and patients with severe hepatic impairment, as in these conditions unbound Linzagolix mean exposure is found increased<sup>3</sup>.

Contraindicated in pregnancy or breastfeeding and patients with osteoporosis or genital bleeding of

unknown aetiology.

### **Current status**

Oral contraceptives and progestogens were effective in almost two-third females suffering from symptomatic leiomyomas. It is clinically known that oestrogen plays an important role in pathogenesis, oral GnRH antagonists may prove effective, especially in non-respondents to progestogens<sup>4</sup>.

Linzagolix is approved for treatment of moderate-to-severe symptoms of leiomyomas in reproductive females in EU (17 June 2022) and in USA under regulatory review and under phase 3 clinical development for management of pain associated with endometriosis<sup>3</sup>.

### **REFERENCES**

1. Mayo clinic. Uterine fibroid [Internet]. [last updated 2022 Sep 21]. Available from: <https://www.mayoclinic.org/diseases-conditions/uterine-fibroids/symptoms-causes/syc-20354288#:~:text=Uterine%20fibroids%20are%20noncancerous%20growths,almost%20never%20develop%20into%20cancer>.
2. Wikipedia. Linzagolix [Internet]. [last updated 2022 Aug 25]. Available from: <https://en.m.wikipedia.org/wiki/Linzagolix>
3. Keam SJ. Linzagolix: First Approval. Drugs. 2022 Aug 23;1-9.
4. Donnez J, Dolmans MM. GnRH antagonists with or without add-back therapy: A new alternative in the management of endometriosis?. *International journal of molecular sciences*. 2021 Oct 20;22(21):11342.



## CASE REPORT

### A Rare Case of Pulmonary Nocardiosis in Late Post Renal Transplant Period

Tushar Gupta\*, Vinay Malhotra\*\*, Seetaram Singh\*, Niranjana Gogoi\*, Dhananjai Agrawal\*\*, Gigin SV\*

#### ABSTRACT

Nocardiosis is an uncommon but important infection in solid organ transplant recipients. It is a life-threatening disease in these patients. We report a case of 44-years-old kidney recipient who developed pulmonary nocardiosis that was successfully treated with intravenous imipenem and doxycycline in conjunction with a reduction in immunosuppressive therapy. This case emphasizes the role of new potent immunosuppressants and diabetes in the occurrence of opportunistic infections. In transplant recipients, who present with pulmonary symptoms and do not respond to usual antibiotics, a Nocardial infection should be suspected.

**Key words:** Immunosuppressive therapy; Pulmonary nocardiosis; Renal transplantation.

#### INTRODUCTION

Nocardiosis is a lethal disease in solid organ transplant recipients. Nocardiosis belongs to the order Actinomycetales. It is a weakly acid fast, Gram-positive, branching filamentous aerobic bacteria. The most frequent species associated with infections in humans are *Nocardia asteroides*, *Nocardia brasiliensis*, *Nocardia farcinica* and *Nocardia nova*.

The usual mode of acquiring infection is through inhalation resulting in a pneumonitis followed by a dissemination of the infection, or can occur through skin trauma. Its three clinical forms include: cutaneous, pulmonary and disseminated. The cutaneous form is related to the transcutaneous inoculation and is common in tropical and warm temperate areas. The most common mode of transmission remains inhalation, resulting in pulmonary localization.

*Nocardia* is an opportunistic pathogen, causing

pulmonary and systemic infections in immunocompromised patients. Patients in an immunocompromised state and having disseminated forms have poor prognosis.

In India, nocardiosis was reported in 1.4% of renal transplant recipients<sup>1</sup>, most common species being *Nocardia asteroides*. Intense immunosuppression is the commonest predisposing factor.

We report herein a rare case of pulmonary nocardiosis developing in late post renal transplant period.

#### CASE REPORT

A 44 years old male patient, with end stage renal disease secondary to an unknown nephropathy, received renal transplantation in November 2019. The donor was his wife. They had 3 HLA mismatch. He had been treated for Hepatitis C before renal transplantation. He was given induction with Basiliximab and started on triple immunosuppressive therapy of Prednisolone, Tacrolimus and Mycophenolate mofetil. He had delayed graft function with serum creatinine being 4.1 mg/dl on postoperative day 7. Graft biopsy done, was reported as mixed rejection with acute tubular necrosis. He received methylprednisolone pulse therapy, intravenous immunoglobulin, plasmapheresis-4 sessions, 3 doses of anti-thymocyte globulin. Patient was discharged with good and stable renal function (serum creatinine of 1.5 mg/dl) on 15<sup>th</sup> postoperative day. New onset diabetes mellitus occurred fifteen months later and was managed by oral anti diabetic drugs.

In August 2021 (21 months post renal transplantation), he was admitted to the Nephrology department with complaints of fever, left sided pleuritic

---

\*Resident, \*\*Senior Professor, Department of Nephrology, SMS Medical College, Jaipur

#### Corresponding Author:

Dr. Tushar Gupta, DM Resident

C-32, Street No.3, North Chajjupur, 100 Feet Road, Shahdara, Delhi-110094

E-mail address: [drtushargupta90@gmail.com](mailto:drtushargupta90@gmail.com)

Mobile No.: 9899907468

chest pain, shortness of breath along with cough and expectoration for 15 days.

Clinical examination revealed: body temperature of 39.6°C, respiratory rate of 28 per minute, blood pressure was 80/60 mm Hg, oxygen saturation of 82% on room air and 97% with 6 L/min oxygen support. Lung examination revealed tachypnoea and crepitations along left lung areas.

Laboratory investigations revealed haemoglobin of 14.8 m/dl, a total leucocyte count of 20,100/mm<sup>3</sup>, neutrophils 90%, lymphocytes 6% and ESR was 58 mm in the first hour. Random blood sugar was 460 mg/dl. Renal function showed blood urea of 80 mg/dl and serum creatinine of 2.0 mg/dl. Other blood investigations which includes serum electrolytes and liver function tests were normal.

Chest radiograph showed non-homogenous opacity in left upper and lower zone [Figure 1]. CT chest revealed consolidation with internal air bronchogram, ground glass haziness in left upper and lower lobe suggestive of pneumonitis [Figure 2].

Microbiological investigations included sputum Gram stain which showed the presence of slender, Gram-positive, branching filamentous bacilli.

The patient was started on higher antibiotics, insulin infusion for blood sugar control and inotropic support for maintaining blood pressure. Sputum sent for modified Ziehl-Neelsen staining, revealed numerous acid-fast branching filamentous organisms, morphologically consistent with *Nocardia* species. [Figure 3].

Trimethoprim-sulfamethoxazole at therapeutic dose was added along with intravenous imipenem-based treatment and oral doxycycline. Moreover, immunosuppressant drugs were temporarily reduced by decreasing dose of tacrolimus by 50% and stopping mycophenolate mofetil. Unfortunately, on treatment, the patient's condition got deteriorated and he succumbed to the illness.

## **DISCUSSION**

Nocardiosis is an infrequent infection and its diagnosis is difficult. Infection occurs in severely immunocompromised patient with reduced-cellular mediated immunity such as solid organ transplants, human immunodeficiency virus infected patients, autoimmune diseases, neoplasia and chronic lung disease. The

most common risk factors are corticosteroid therapy and immunosuppression.

An observation by Carnet et al concluded that patients having Tacrolimus based immunosuppression were at increased risk for nocardiosis when compared to those on cyclosporin<sup>2</sup>.

Our patient received induction treatment with Basiliximab and was maintained on mycophenolatemofetil, steroids, and Tacrolimus. He also received intravenous immunoglobulin, plasmapheresis, anti-thymocyte globulin for treatment of acute rejection. Moreover, diabetes mellitus is an additional risk factor favouring occurrence of opportunistic infection in our patient.

Nocardial infection is most common in first six months' post-transplant period<sup>3</sup>. Rarely reported after the first year of transplantation<sup>4</sup>, but in our case presented after twenty-one month<sup>5</sup> post-transplant.

In pulmonary nocardiosis, the most common clinical presentation is a sub-acute or chronic necrotizing pneumonia. Pulmonary nocardiosis has a quintessential presentation, which includes fever, malaise, cough, anorexia, dyspnea, and chest pain. Our patient also had fever, cough and breathlessness as presenting complaints.

The most important diagnostic tool of pulmonary nocardiosis is Gram staining of sputum. However, no specific clinical or radiological features suggest nocardiosis. Only one third of sputum samples may show *Nocardia*, so multiple sputum specimens have to be examined.

Case studies showed that about 60-80% of nocardiosis in renal transplant patients is pulmonary nocardiosis and half of them have isolated lung involvement. A solitary pulmonary involvement was present in our patient without clinically apparent dissemination.

The most affected organs include: the lungs, skin, subcutaneous tissue and cerebro-nervous system. Other locations have been described: cardiac, ocular, osteoarticular.

So, for management of pulmonary nocardial infections, the key is to have a high index of suspicion, early diagnosis and adequate treatment. In transplant recipients, who present with pulmonary symptoms and do not respond to usual antibiotics, a Nocardial infection should be suspected.

Radiological patterns of pulmonary nocardiosis include, non-specific findings such as air space consolidation, presence of irregular nodular lesions, which may or may not associated with cavitation. Mediastinal and hilar lymphadenopathy may also be seen. In our patient, multilobar involvement was present with airspace consolidation but cavitation was lacking.

Prophylaxis with cotrimoxazole is recommended in renal transplant patients for pneumocystis carinii and urinary tract infection, and it is believed by some authors to protect from Nocardia infection. However, there is increasing cases of breakthrough infections in patients taking TMP-SMX prophylaxis.

Sulphonamides, mainly cotrimoxazole, are the treatment of choice, although other drugs, such as imipenem, amikacin, linezolid, cefotaxim, clarithromycin, ofloxacin, amoxicillin-clavulanic acid and tetracycline derivatives are safe and effective.

Usually, six to nine months of antibiotic therapy, in case of localized pulmonary infections are needed and nine to twelve months in case of cerebro nervous system involvement.

Reduction of immunosuppression may be a helpful adjunctive therapy in severe forms of the disease but it is not a mandatory approach.

In pulmonary localization, mortality is about 40% and increases to 64% in disseminated nocardiosis and 100% in the presence of cerebro nervous system involvement<sup>5</sup>.

## CONCLUSION

Transplant physicians should be aware of this rare infection and consider nocardiosis in differential diagnosis of pneumonia even in late post-transplant period, especially when the radiological features are atypical and in patients who have not responded to the empirical treatment.



Figure 1. Chest X-ray




Figure 2. CT chest




Figure 3. Modified ZiehlNeelsen stain

## REFERENCES

1. John GT, Shankar V, Abraham AM, Mathews MS, Thomas RP, Jacob CK. Nocardiosis in tropical renal transplant recipients. Clin Transplant 2002; 16:285-9.
2. Canet S, Garrigue V, Bismuth J, Chong G, Lesnik A, et al. (2004) Nocardiosis- is it frequently observed after the introduction of new immunosuppressive agents in renal transplantation? Nephrologie 25: 43-8.
3. Fishman JA, Rubin RH. Infection in organ transplant recipients. N Engl J Med 1998;338:1741-51.
4. Yu X, Han F, Wu J, He Q, Peng W, Wang Y, et al. Nocardia infection in kidney transplant recipients: Case report and analysis of 66 published cases. Transpl Infect Dis 2011;13:385-91.
5. Martinez Tomas R, Menendez Villanueva R, Reyes Calzada S, Santos Durantes M, Valles Tarazona JM, et al. (2007) Pulmonary nocardiosis: risk factors and outcomes. Respirology 12: 394-400.

### Vaccination for Elderly: Easiest Way for Disease Prevention

Lakshmi Kant Goyal\*, Sunny Singhal\*\*, Monica Jain\*\*\*, Shashank Sharma\*\*\*\*

The health problems faced by the elderly are unique compared to other age groups. In elderly, immune function are decreased predisposing this population prone for infectious diseases. The co-morbid conditions including chronic lung diseases and diabetes etc. make the infections more severe. The recovery from infections is also slow and usually takes longer time in elderly. In India, preventive care for diseases is often ignored among elderly (specially for age above 65 years). Many infections and illnesses could be easily avoided by administration of the right vaccines and diagnostic health screenings. Sadly, we don't have a National Immunization Schedule for the elderly as we have for children. It is expected from the family members to take proactive steps to vaccinate their older adults in the family to keep family stay healthy. Following is a brief note about preventive vaccination for older adults.

Elderly are more susceptible to the pneumococcal disease, influenza, herpes zoster, tetanus and COVID-19.

Pneumonia and other pneumococcal related diseases and their related complications are more common and severe in elderly compared to younger ones. For these people, pneumococcal vaccine is of two types: the 23-valent pneumococcal polysaccharide vaccine (PPSV23), and the 13-valent pneumococcal conjugate vaccine (PCV13). The PCV13 is more immunogenic. PPSV23 is less immunogenic but covers 10 more strains than PCV13. For our seniors, the recommended schedule is 1 dose PCV13 if previously did not receive PCV13, followed by 1 dose PPSV23 at least 1 year after PCV13. If previously received PPSV23 but not PCV13 at age 65 years or older; 1 dose PCV13 at least 1 year after PPSV23 is recommended. PCV13 and PPSV23 should not be

administered during same visit. The recommended interval between PCV13 and PPSV23 is  $\geq 1$  year for immunocompetent elder. However, for those with immunocompromising or special conditions (chronic renal failure, nephrotic syndrome, blood cancer, generalized malignancy, solid organ transplant, multiple myeloma or splenectomy), the recommended time to wait is  $\geq 8$  weeks.

Influenza (flu) related morbidity and mortality is also higher in elderly. These people should receive 1 dose of inactivated influenza vaccine or a recombinant influenza vaccine annually preferably before the onset of influenza activity in the community, preferably in October before the flu season starts in India. Elderly should get a flu shot and not a nasal spray vaccine.

Herpes Zoster or Shingles, occurs when the chickenpox virus (dormant in almost all adults who had chickenpox in childhood); reactivates in later life. The condition often brings a blistering, painful rash. Post herpetic pain is more common and severe in elderly. There are 2 vaccines for shingles prevention. A live attenuated zoster vaccine (1-dose) and an adjuvant herpes zoster subunit vaccine (2 doses, administered 2 to 6 months apart, minimum 4 weeks interval). The adjuvanted herpes zoster subunit vaccine (2 dose vaccine) has higher efficacy. If the person had previously vaccinated with live attenuated zoster vaccine, then adjuvant herpes zoster subunit vaccine can be administered at least 2 months after. Live vaccines are contraindicated in severe immunocompromising conditions.

Diphtheria toxoid, tetanus toxoid and acellular pertussis-containing vaccines help in protecting against diphtheria, tetanus and pertussis, but they will not prevent

---

\*Associate Professor, Department of Geriatric Medicine

\*\*Assistant professor, Department of Geriatric Medicine

\*\*\*Senior Professor, Department of Pharmacology

\*\*\*\*Senior Demonstrator, Department of Forensic Medicine

#### Corresponding Author:

Dr. Lakshmi Kant Goyal

Associate Professor, Department of Geriatric Medicine

Mob. : 9462651019 Email : drlkgoyal@gmail.com



## Vaccination for Elderly: Easiest Way for Disease Prevention

all cases. Tdap (Tetanus and diphtheria toxoids and acellular pertussis vaccine), Td (Tetanus and diphtheria toxoids) and TT (Tetanus toxoid) are available. The elderly people should receive 1 dose Tdap, then Td or Tdap booster every 10 years. Previously TT was used as booster. COVID-19 (CORONA) related pneumonia occurs in more

severe form in elderly persons. Vaccination for COVID-19 is also available and recommended for our seniors.

Apart from these vaccines for elderly, there are several other vaccines available i.e. Hepatitis B vaccine, which can be given according to the individual risk of getting infections.

### Recommendation of vaccination in persons aged $\geq 65$ years

Pneumococcal vaccine	Not previously vaccinated	A dose of PCV13 followed by a dose of PPSV23, 1 year after
	Previously vaccinated with only PPSV23	A dose of PCV13, one year after vaccination with PPSV23
	PPSV23 received at age $< 65$ years	PCV13 at age $\geq 65$ years and PPSV23 after 1 year (at least 5 years after previous dose of PPSV23)
Influenza (flu)		Every year
Herpes Zoster	Adjuvant herpes zoster subunit vaccine (RZV)	2 doses, 2 to 6 months apart, minimum 4 weeks interval
	A live attenuated zoster vaccine (ZVL)	1 dose
Tetanus Toxoid	Tdap (Tetanus and diphtheria toxoids and acellular pertussis vaccine), Td (Tetanus and diphtheria toxoids)	1 dose Tdap, then Td or Tdap booster every 10 years
COVID-19 (CORONA) vaccine	Available	



# The Rajasthan Medical Journal

## Guidelines / Instructions

Articles are accepted on the basis of significance, scientific merit and applicability. Authors are requested to make their report on the basis of original research carried out by themselves or their groups. The manuscripts will be reviewed for possible publication with the understanding that they are being submitted only to The Rajasthan Medical Journal and have not been published, simultaneously submitted, or already accepted for publication elsewhere. The editors review all submitted manuscripts initially and reject outright manuscripts that do not carry a clear message to the intended readership or are of insufficient originality.

The RMJ follows a policy of blind, peer review without revealing the identity of the contributors and each manuscript typically is read by at least two or three reviewers with expertise in the study's subject matter and methodology. All comments by reviewers are confidential and shall not be published. Within a period of around twelve weeks, the contributors will be informed about the reviewer's comments and acceptance or rejection of the manuscript revisions, if any, suggested by the editors or reviewers are communicated to the author's online requesting submission of the revised manuscript. The editor will make every effort to insure that authors informed as to the disposition of their manuscript within three months of receipt of the submission. The RMJ reserves the right to copy-edit the articles for length, grammar, punctuation, print style and format. Manuscripts must confirm to the guidelines for avoiding ethnic biases and sexist language. Research involved by an Institutional review board.

## Online Submission of Manuscripts

Authors should submit their manuscript to The RMJ through its website [www.rmjonline.in](http://www.rmjonline.in) using the link "Submit Article". Electronic submission substantially reduces the editorial processing and peer reviewing times and shortens overall publication time.

**Note:** If you encounter any difficulty in submitting the manuscripts through the website link, you can e-mail all the manuscripts to the Editor as an attachment at [editor@rmjonline.in](mailto:editor@rmjonline.in)

For online submission, articles should be prepared in two files (First Page file and The Article file). Images should also be attached with these two files.

**First Page File:** Prepare the title page, covering letter, acknowledgement, etc. using a word processor program. All

information which can reveal your identity should be here. Use text/doc/rtf/doc/PDF files. Do not zip the files.

**The Article File:** The main text of the article beginning from abstract till references (including tables) should be in this file. Do not include any other information (acknowledgment, your name in page headers, etc.) in this file. Use text /rtf/doc/PDF files. Do not zip the files. Limit the file size to 400 KB. Do not incorporate images in the file. If the file size is large, graphs can be submitted as images separately without incorporating them in the article file so as to reduce the size of the file.

**Images:** Submit good quality color images please! Each image should be less than 100 KB in size. Size of the image can be reduced by decreasing the actual height and width of the image (keep up to 400 pixels or 3 inches). JPEG image format is the most acceptable. Do not zip the files.

**Legends:** For the figures/images should be kept ready for copy paste during the submission process. Once a manuscript is submitted online, please send hard copies of the image and Contributors' Form to the Editor's address within four weeks. A Compact Disk or a Pen Drive containing the manuscript should be avoided.

## Instructions for authors regarding manuscript submission

- Before you submit your manuscript, please read the editorial policy and the editorial guidelines of The RMJ carefully and make sure that your manuscript confirms to them. Manuscripts may be returned to the authors if they violate The RMJ editorial policy or deviate from its editorial guidelines.
- In order to widen the scope of communication, the following formats are made available - Editorials, review articles, original papers, leading articles, post graduate clinical updates, general practitioner updates, clinico-pathological conferences, unusual cases - case reports, clinical puzzles, readers corner and so on.

## Physical Submission of Manuscripts

Send three copies of the manuscript, three sets of figures along with a covering letter, Contributors Form duly signed by all the contributors, checklist and a compact disk. Place the figures in a separate envelope. The covering letter must include information on prior or duplicate publication or submission elsewhere of any part of the work/study; and a statement of financial or other relationships that might lead to a conflict of interest. Copies of any permission(s) to reproduce published material, and to use illustrations or report information about identifiable people must accompany the manuscript. Send all the material to:



# The Rajasthan Medical Journal

The Editor,

The Rajasthan Medical Journal (RMJ),

S.M.S. Medical College,

J.L.N. Marg, Jaipur – 302 004, Rajasthan, INDIA

It is essential that you provide complete contact details including full institutional address, telephone numbers, complete postal address and e-mail address for the corresponding author.

## **Type of Manuscripts and Word Limits**

**Leading Articles:** The editors commission leading articles that are 1,000-1,500 words in length and address topics of current interest. They should be accompanied by no more than 10 references. Submissions are subjected to editorial review before acceptance. The editor retains the right to alter style and shorten material for publication.

**Review Articles:** The editorial board encourages submission of review articles on topics of current interest. The manuscript should be restricted to 4,000 words and up to 50 references. An abstract of no more than 250 words and up to 6 keywords should be provided.

**Original Papers:** These should be in the format of a) Introduction, b) Patients and methods, c) Results, and d) Discussion. The manuscript should be restricted to 3,000 words, up to 30 references and carry no more than 4 tables or figures. Please provide a 250 words structured abstract and up to 6 keywords. Structured abstract should include - background, study objective, design, setting, sample/population, duration of study, results and conclusion. Below the abstract, provide up to 6 keywords that will assist cross-indexing the articles and may be published with the abstract.

**Unusual Cases:** This section presents reports on rare cases. The manuscript should be in the format of a) Introduction, b) Case report, and c) Discussion. Submissions to this section should carry no more than 1,500 words, 2 figures and 5 references. An unstructured abstract of up to 150 words and 6 keywords should be provided.

**Reader's Corner:** Comments on papers recently published in the Journal. The letters should be restricted to up to 500 words and 3 references and should not carry figures.

**Post Graduate Clinical Update:** This category is for a concise, comprehensive summary of a topic of interest for post graduate students. The word count should not exceed 4,000

words.

**Clinical Puzzle:** This category is for unusual cases that makes an education point. Since the aim of these articles is to stimulate the reader to think about the case, the title should be ambiguous and not give away the final diagnosis immediately. Clinical puzzle should be in two parts. The first part should contain a brief description of the case (maximum 250 words) followed by an image and question-designed as task 1, 2, 3, 4, and so no. The legend of the image should not indicate the diagnosis but should simply describe the nature of the image. Each task is to be followed by a short description of what actually was done in the specific case. The answers to all the tasks should come in the second part, followed by a brief key message indicating few learning points. Clinical puzzle will not include more than 5 references, a maximum of 2 images, 2 figures and an author limit of 5. The quality of the image should be no less than 600 dpi and should be in JPG format. We do ask that authors indicate that they have obtained patient consent.

**General Practitioner Update:** This category presents a simple concise and clear summary of a topic of interest to general practitioners. The word count should not exceed 4,000 words. The word count excludes the title page, abstract, tables, acknowledgements & contributors, and the references.

## **Preparation of the Manuscript**

Send laser print-out on thick paper of A4 size (8.27" x 11.69"), with margins of 1 inch on all side. Print on only one side of the paper. Use double spacing throughout. Number the pages consecutively, beginning with the title page. The language should be American English.

**Title Page:** Should carry the following information:

- Title of the article
- Running title or short title of no more than 50 characters
- Name of each contributor (Last name, First Name and initial of Middle Name)
- Name of the department(s) and institution(s) to which the work should be attributed
- Name, address, phone number, facsimile number and e-mail address of the contributor responsible for correspondence
- Type of manuscript (Original/Review/Unusual Cases, etc.)
- Word counts separately for abstract and for the text



# The Rajasthan Medical Journal

(excluding the references and abstract)

**Acknowledgement:** Specify contributions that need an acknowledgement but do not justify authorship, such as general support by a departmental head or acknowledgements of technical, financial and material support and if the manuscript was presented as a part at the meeting in the organization, place and exact date on which it was read.

**Abstract Page:** The second page should carry the full title of the manuscript and an abstract (see above for word limits). For original articles, the abstract should be structured and arranged in the format indicated earlier.

## **Text of the Article:**

- State the purpose of the article and summarize the rationale for the study or observation in the introduction. For unusual cases, give incidence of similar cases in the past.
- Describe the selection of the observational or experimental subjects clearly in patients and methods section. Identify the age, sex and other important characteristics of the subject. Identify the methods, apparatus (give the manufacturer's name in parentheses) and procedures in sufficient detail.
- Give references to established method, describe new or substantially modified methods, give reasons for using them, and evaluate their limitation.
- Identify precisely all drugs and chemicals used, including generic name(s) dose(s) and route(s) of admission.
- Report of randomized clinical trials should be based on the consort statement.
- Do not use patients' names, initials or hospital numbers especially in the illustrative material.
- Present the results in logical sequences in the text, tables and illustrations; emphasize or summarize only important observations. Use standard guidelines for statistics.
- Emphasize the new and important aspects of the study and the conclusion that follow from them along with implications of the findings and their limitation in the discussion section.

## **References:**

- References should be numbered consecutively in the order in which they are first mentioned in the text.
- Identify references in text, tables, and legends by Arabic

numbers in superscript. References cited only in tables or figure legends should be numbered in accordance with the sequence established by the first identification in the text of the particular table or figure.

- The titles of journals should be abbreviated according to the style used in Index Medicus.
- Avoid using abstracts, un-published observations and personal communications as references.

## **Tables:**

- Table should be self-explanatory and should not duplicate textual material.
- Tables with more than 10 columns and 25 rows are not acceptable. Limit the number to minimum required. Number tables in Arabic numerals, consecutively in the order of their first citation in the text and supply a brief title for each.
- Place explanatory matter in footnotes, not in the heading. Explain in footnotes all non-standard abbreviations that are used in each table. For footnotes, use the following symbols in the sequence: \*, #, \$, \*\*.
- Obtain permission for all fully borrowed, adapted and modified tables and provide a credit line in the footnote.

## **Illustrations (Figures):**

- Submit three sets of sharp, glossy, un-mounted, color photographic prints, with height of 4 inches and width of 6 inches. Computerized color print-outs are not acceptable.
- Figure should be numbered consecutively according to the order in which they have been first cited in the text.
- Each figure should have a label pasted on its back indicating the number of the figure, the running title, top of the figure and the legends of the figure. Do not write on the back of figures, scratch, or mark them by using paper clips.
- Symbols, arrows or letters used in photomicrographs should contrast with the background and should be marked neatly with the background and should be marked neatly with transfer type or by tissue overlay and not by pen.
- If a figure has been published, acknowledge the original source and submit written permission from the copyright holder to reproduce the material. A credit line should appear in the legend for such figures.
- Type or print out legends (maximum 40 words,





# The Rajasthan Medical Journal

excluding the credit line) for illustrations with Arabic numerals corresponding to the illustrations. When symbols, arrows or letters are used to identify parts of the illustration, identify and explain each one clearly in the legend explain the internal scale and identify the method of staining in photomicrographs.

## Submitting a Revised Manuscript

While submitting a revised manuscript, contributors are requested to include, along with a single copy of the final revised manuscript, a photocopy of the revised manuscript with the changes underlined in red and with the point to point clarification to each comment. The manuscript number should be written on each of these documents.

## Reprints

Journal does not provide any free printed reprints. Reprints can be purchased at the time of submitting the proofs.

Checklist (To be tick marked as applicable and one copy attached with the manuscript)

- Covering letter duly signed by all the contributors
- Previous publications / presentations mentioned
- Source of funding mentioned
- Conflicts of interest disclosed
- Authors
- a. Middle Name initials provided
- b. Author for correspondence, with e-mail address provided
- c. Identity not revealed in paper except title page (e.g. Name of the institute in material and methods, citing previous study as 'our study', name of institute in photographs, etc.)
- Presentation and format
  - a. Double spacing
  - b. Margins 2.5 cm from all four sides
- c. Title page contains all the desired information (vide supra)
- d. Running title provided (not more than 50 characters)
- e. Abstract page contains the full title of the manuscript
- f. Abstract provided (150 words for case reports and 250 words for original articles)
- g. Structured abstract provided for an original article
- h. Keywords provided (three or more)
- i. Headings in title case (not ALL CAPITALS, not underlined)
- j. Reference cited in superscript in the text without brackets
- k. Reference according to the journal's instructions
  - Language and Grammar
    - a. Uniformly American English
    - b. Abbreviations spelt out in full for the first time
  - Tables and Figures
    - a. No repetition of data in tables/graphs and in the text
    - b. Actual numbers from which graphs drawn to be provided
    - c. Figures necessary and of good quality (color)
    - d. Labels pasted on back of the figures (no names written)
    - e. Figure legends provided (not more than 40 words)
    - f. Patients' privacy maintained (if not, written permission enclosed)
    - g. Table and figure numbers in Arabic letter (not Roman)
    - h. Credit note for borrowed figures / tables provided
  - Manuscript provided on a Compact Disk



# The Rajasthan Medical Journal

Contributors Form (to be modified as applicable and one single copy attached with the manuscript)

Title: \_\_\_\_\_

Manuscript Submission date \_\_\_\_\_ (for manuscripts submitted online) I / we certify that I / we have participated sufficiently in the intellectual content, conception and design of this work or the analysis and interpretation of the data (when applicable), as well as the writing of the manuscript, to take public responsibility for it and have agreed to have my / our name listed as a contributor. I / we believe the manuscript presents valid work. Neither this manuscript nor one substantially similar content under my / our authorship has been published or is being considered for publication elsewhere, except as described in the covering letter. I / we certify that all the data collated during the study is presented in this manuscript and no data from the study has been or will be published separately. I / we attest that, if requested by the editors. I / we will provide the data // information or will cooperate fully in obtaining and providing the data/information on which the manuscript is based, for examination by the editors or their assignees. I / we also certify that we have taken all necessary permissions from our institution and / or department for conduction and publishing the present work. Financial interests, direct or indirect, that exist or may be perceived to exist for individual contributors in connection with the content of this paper have been disclosed in the cover letter. Sources of outside support of the project are named in the cover letter.

I / we hereby transfer (s), assign(s), or otherwise convey(s) all copyright ownership, including and all rights incidental there to, exclusively to The Rajasthan Medical Journal (The RMJ) in the event that such work is published by The RMJ. The RMJ shall own the work, including 1) copyright; 2) the right to grant permission to republish the article in whole or in part, with or without fee; 3) the right to produce preprints or reprints and translate into languages other than English for sale or free distribution; and 4) the right to republish the work in a collection of articles in any other mechanical or electronic format. We give the rights to the corresponding author to make necessary changes as per the request of the journal, do the rest of the correspondence on our behalf and he/she act as the guarantor for the manuscript on our behalf. All persons who have made substantial contributions to the work reported in the manuscript, but who are not contributors, are named in the acknowledgment and have given me/us their written permission to be named. If I/we do not included an acknowledgment that names I / we have not received substantial contributions from non-contributors and no contributor has been omitted.

	Name	Signature	Date Signed
1	_____	_____	_____
2	_____	_____	_____
3	_____	_____	_____
4	_____	_____	_____
5	_____	_____	_____
6	_____	_____	_____



# The Rajasthan Medical Journal

N. B. If you have already renewed your subscription for 2020, please disregard this notice

Editor

The Rajasthan Medical Journal

Name \_\_\_\_\_ Number of Copies \_\_\_\_\_

Organization (if applicable) \_\_\_\_\_ Department: \_\_\_\_\_

Position: \_\_\_\_\_ Designation \_\_\_\_\_

Address: \_\_\_\_\_

Contact No. \_\_\_\_\_ Fax \_\_\_\_\_

Email \_\_\_\_\_

Please circle the rate you are wanting

- |                            |          |
|----------------------------|----------|
| 1. Annual Membership Fee - | Rs. 500  |
| 2. Life Membership Fee –   | Rs. 2000 |

Thank you for renewing your details. The RMJ will send you an invoice for 2017

Please find enclosed cash / Cheque / DD No. \_\_\_\_\_ Dated \_\_\_\_\_

Drawn on (Bank Name) \_\_\_\_\_

For Rs. \_\_\_\_\_ favoring the Principal, SMS Medical College, Jaipur

Please invoice ☐

New Subscription ☐

Renewal ☐

